

Ginstar® EC COTTON DEFOLIANT



Supplemental Label

For Agricultural Use Only

ACTIVE INGREDIENTS: Thidiazuron: N-phenyl-N'-1,2,3-thiazol-5-ylurea*	12%
Diuron: 3-(3,4-dichlorophenyl)-1,1-dimethylurea*	6%
INERT INGREDIENTS:	82%
Contains 1 lb. Thidiazuron per gallon <i>and</i> 0.5 lb. Diuron per gallon.	TOTAL 100%

*Protected by U.S. Patent Nos. 4,294,605 and 4,613,354.

EPA Reg No. 264-634

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN DANGER — PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-888-AVENTIS (1-888-283-6847)

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Hold eyelids open and immediately flush with a steady, gentle stream of water for at least 15 minutes. Call a physician.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate use of gastric lavage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive. Causes irreversible eye damage. Causes burns. Harmful if inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing dust or spray mist. Prolonged or frequently repeated skin-contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers (other than mixers and loaders) must wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber gloves
- Chemical-resistant footwear plus socks
- Protective eyewear

Mixers and loaders must wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate or butyl rubber gloves
- Chemical-resistant footwear plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4–6)], the handler PPE requirements may be reduced or modified as specified in the WPS

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Do not apply when weather conditions favor drift from the target area.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container and keep closed. Store in a cool, dry place. Do not use or store near heat or open flame. Do not contaminate food or foodstuffs.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in sanitary landfill, or by other procedures approved by State and local authorities.

DO NOT REUSE EMPTY CONTAINER

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

In case of spillage, cover with an absorbent such as soda ash, lime, clay, or sawdust. Sweep up and bury. Wash area thoroughly with detergent and water.

GENERAL INFORMATION

GINSTAR® EC is a defoliant to be used as a pre-harvest aid for cotton. GINSTAR® EC Cotton Defoliant has performed well under both cool and warm weather conditions.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system.

Rainfall within 12 hours after application can reduce the effectiveness of GINSTAR® EC Cotton Defoliant.

Do not feed foliage from treated cotton plants or gin trash to livestock.

Do not plant the following crops earlier than the specified periods after application of GINSTAR® EC:

small grains, sorghum, corn, or root crops (except carrots, onions).....	two (2) months
legumes (including alfalfa) or leafy vegetables (except lettuce)	two (2) months
cole crops, garlic, safflower, tomatoes, and watermelon	two (2) months
carrots	three (3) months
onions	four (4) months
cantaloupe, honeydew melon/ casaba melon, muskmelon, or peppers	five (5) months
lettuce	two (2) months
	<i>with deep-plowing of soil (12–15 inches);</i>
	<i>or nine (9) months when soil is only disced (4–6 inches).</i>

Cover Crops: Small grains and/or legumes intercropped within the cotton crop to which GINSTAR® EC Cotton Defoliant will be applied may only be used as cover crops and may not be harvested for food or feed. Small grains and/or legumes planted earlier than two (2) months following GINSTAR® EC application may only be used as cover crops and may not be harvested for food or feed.

Do not plant any other rotational crops (*except* those specified above) within one year of application of GINSTAR® EC.

Do not use immature crops for food or feed.

Do not allow spray drift to contact trees or crops other than the target crop of mature cotton, or cotton you desire to defoliate, as this product may injure or defoliate other crops.

Particular care should be taken when applying GINSTAR® EC adjacent to lettuce, citrus, or cantaloupe.

PRACTICES TO LOWER THE POTENTIAL FOR DRIFT ONTO NONTARGET CROPS

During applications, particularly under windy conditions, GINSTAR® EC Cotton Defoliant may drift to nontarget crops. To help reduce the drift potential, use the following practices:

- Do not apply GINSTAR® EC by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20–30 psi) is recommended.
- Do not apply GINSTAR® EC when a temperature inversion is present or when conditions favor an inversion prior to completing application(s).
- Additionally, do not apply GINSTAR® EC by air within one-half (1/2) mile of lettuce. Do not apply GINSTAR® EC by ground equipment within 100 feet of lettuce.

PRACTICES TO LOWER DRIFT POTENTIAL ONTO NONTARGET CROPS IN THE RIO GRANDE VALLEY OF TEXAS

To help reduce the drift potential, use the following practices:

- Do not apply GINSTAR® EC Cotton Defoliant by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20–30 psi) is recommended.
- Do not apply GINSTAR® EC when a temperature inversion is present or when conditions favor an inversion prior to completing application(s).
- Use the largest nozzle orifice possible which permits proper deposition and coverage of product.
- In addition, for citrus crops, do not apply GINSTAR® EC by air when citrus in flush is within five (5) miles downwind of the point of application. Do not apply GINSTAR® EC by ground when citrus in flush is within one-half (1/2) mile downwind of the point of application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate or butyl rubber gloves
- Chemical-resistant footwear plus socks
- Protective eyewear

TIME OF APPLICATION

Apply GINSTAR® EC Cotton Defoliant only to mature cotton plants when the last boll you expect to harvest is mature. A boll can be described as "mature" when it is too hard to be dented when squeezed between thumb and fingers, is difficult to slice with a sharp knife, and/or when seeds cut in cross sections have fully developed cotyledons, as evidenced by an absence of jelly within the seed.

Apply GINSTAR® EC at least 5 days prior to anticipated harvest.

PRECAUTION: The addition of adjuvants can cause desiccation and/or leaf freezing during periods of high temperature. The use of compounds that desiccate leaf tissue is not recommended.

Use of GINSTAR® EC Cotton Defoliant under extremely cool or adverse conditions can result in less than desirable defoliation and/or growth inhibition.

APPLICATION

GINSTAR® EC Cotton Defoliant may be applied by air or ground equipment. Apply specified dosages in 10–25 gallons of spray per acre with ground equipment or 2–10 gallons per acre by aircraft.

DOSAGE

Apply GINSTAR® EC Cotton Defoliant at a rate of 0.4 to 1.0 pint of formulated product per acre prior to harvest (see *Application Rate Table*). At some locations, following the initial GINSTAR® EC application, it may be necessary to make a second application of GINSTAR® EC (DO NOT exceed 1.0 pint/acre per season), or an application of a standard defoliant.

MIXING INSTRUCTIONS

Fill the spray tank with one-half of the required amount of water. Add the proper amount of GINSTAR® EC Cotton Defoliant and start the spray tank agitator. Finish filling the tank with the balance of water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

**PRECONDITIONING
FOR USE IN CALIFORNIA ONLY**

GINSTAR® EC Cotton Defoliant may be used as a preconditioner to enhance the activity of a defoliant application. Apply GINSTAR® EC at 4–6 oz. formulated product per acre. Apply in 10–25 gallons of water per acre by ground or 2–10 gallons of water per acre by aerial application. Timing of application is recommended 7–10 days prior to a defoliation application of GINSTAR® EC (see *Application Rate Table*) or the use of another defoliant. Refer to the second product label prior to use for complete recommendations.

GINSTAR® EC COTTON DEFOLIANT APPLICATION RATE TABLE

To Achieve an Application Rate of:	Use This Amount of GINSTAR® EC	At the Indicated Rate, One Gallon of GINSTAR® EC Will Treat:
0.075 lbs. ai/Acre	0.4 pts./A (6.4 oz./A)	20 Acres
0.10 lbs. ai/Acre	0.55 pts./A (8.8 oz./A)	15 Acres
0.15 lbs. ai/Acre	0.8 pts./A (12.8 oz./A)	10 Acres
0.1875 lbs. ai/Acre	1.0 pts./A (16.0 oz./A)	8 Acres

DO NOT APPLY MORE THAN 1.0 PINT OF GINSTAR® EC PER ACRE PER SEASON.

CLEANOUT INSTRUCTIONS

Do not allow the spray solution to dry in the application equipment. Immediately following application, clean all equipment (mix tanks, pumps, transfer lines, application tanks, sumps, booms, nozzles, and all related equipment) thoroughly with cleaner and water. Refer to Aventis CropScience technical information bulletins for tank cleaners which may be helpful in removing fresh residues.

Should small quantities of GINSTAR® EC Cotton Defoliant remain in inadequately cleaned equipment, they may be released during subsequent applications and may cause damage to crops. Aventis CropScience accepts no liability for damage to crops due to inadequately cleaned equipment.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aventis CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: AVENTIS CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aventis CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. AVENTIS CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AVENTIS CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Aventis CropScience USA LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

Ginstar EC Cotton Defoliant

EA80414